

COVID-19 OAP DECISION TOOL

Have you been in ‘close contact’ with someone who has tested positive for COVID-19

Or

Are you living in the same household as, being an intimate partner of, or providing care in a non-healthcare setting for a person with confirmed or suspected COVID-19 infection?

Definition of close contact: within 6 feet of the sick contact for a cumulative total of 15 minutes or more in a 24-hour period *or* exposed to respiratory secretions by being directly coughed or sneezed on.

Yes



Vaccination Status

Have you been boosted?
OR
Completed your primary series of Pfizer or Moderna vaccine within the last 6 months?
OR
Completed your primary series of J&J vaccine within the last 2 months?

Yes →

Get tested 5 days after exposure even if you don't have symptoms and wear a mask around others for 10 days.

No or unsure



No



	Identified Risk	Unknown Risk
Asymptomatic People	<u>Quarantine at home for 5 days after last known infectious exposure.</u> After that, wear a mask around others for five additional days. Avoid contact with people at higher risk for severe illness. Get a test 5 days after last known exposure, if possible.	Self-monitor for symptoms for 10 days. Wear a mask around others, and practice good hand hygiene and social distancing in accord with the recommendations for all members of the community.
Symptomatic People	<u>Get a COVID-19 test as soon as you are able, and quarantine at home for 5 days after last known infectious exposure.</u> Contact your <u>primary care physician (or public health) for guidance.</u> After quarantine, wear a mask around others for five additional days. Avoid contact with people at higher risk for severe illness.	Get a COVID-19 test if possible, and isolate at home until you know the result of your test. Avoid contact with people at higher risk of severe illness while you are symptomatic. Pre-notify and consult your primary health care provider for moderate/severe illness.

Symptomatic = fever (≥ 100.4 F), chills, fatigue, muscle or body aches, congestion or runny nose, sore throat, cough, shortness of breath, nausea or vomiting, diarrhea, headache, or new loss of taste or smell.

In absence of specific guidance from local public health professionals this CDC based tool may be used. Specific recommendations may be modified by local authorities.